

311 E 79 th St. 212-996-6633	16 E 52 nd St. 212-826-3903	68 E 86th St. 212-535-1845	1150 Fifth Ave. 212-369-2490	300 Cadman Plaza 718-280-0020
--	---	-------------------------------	---------------------------------	----------------------------------

**48 Hour Colonoscopy
Preparation Instructions:**

Time	Instructions
<u>1 Week Before Procedure</u>	<ul style="list-style-type: none"> Buy at local pharmacy: <ul style="list-style-type: none"> 2 Gatorade bottles 32oz each (NO red, orange, or purple colors) 4 Dulcolax tablets 5mg each for a total of 20mg 2 separate 10oz bottles of Magnesium Citrate for 20oz total Make sure you have your preparation medication (Golytely) by this time
<u>3-5 Days Before</u>	<ul style="list-style-type: none"> Please have your COVID-19 PCR test completed (No Rapid test)
<u>48 Hours (2 days) Before Procedure</u>	<ul style="list-style-type: none"> NO solid foods, only clear liquids including: <ul style="list-style-type: none"> Water Apple, White Grape, & White Cranberry juices without pulp Clear soup Broth Tea or Coffee (No milk, cream) Gatorade / Powerade (NO red, orange, or purple colors) Jell-O (NO red, orange, or purple colors) Popsicles or Sorbet (NO red, orange, or purple colors) No Alcohol
<u>2 Days Before Procedure at 12 Noon</u>	<ul style="list-style-type: none"> Take all 4 Dulcolax tablets with the first 32oz Gatorade bottle (NO red, orange, or purple colors)
<u>2 Days Before Procedure at 3pm</u>	<ul style="list-style-type: none"> Drink 1 of the 10oz bottles of Magnesium Citrate
<u>2 Days Before Procedure at 5pm</u>	<ul style="list-style-type: none"> Drink second 10oz bottle of Magnesium Citrate After, slowly drink the second 32oz Gatorade bottle before the end of the day

Instructions continued on page 2



48 Hour Colonoscopy
Preparation Instructions Continued:

<u>24 Hours Before Procedure</u>	<ul style="list-style-type: none"> • Prepare Golytely <ul style="list-style-type: none"> • Fill container to “fill” line with water and mix until dissolved • Chill if desired • <u>NO Solid food Only Clear Liquid</u> including: <ul style="list-style-type: none"> • Water • Apple, White Grape, & White Cranberry juices without pulp • Clear soup Broth • Tea or Coffee (No milk, cream) • Gatorade / Powerade (No Red, Orange, or Purple Colors) • Jell-O (No Red, Orange, or Purple Colors) • Popsicles or Sorbet (No Red, Orange, or Purple Colors) • NO Alcohol
<u>5pm Night Before Procedure</u> <u>Dose 1</u>	<ul style="list-style-type: none"> • Drink the first half of the solution steadily over the next hour and a half <ul style="list-style-type: none"> • Drink 1 cup (8oz) every 10 minutes or so for a total of 8 cups (64oz) • Half of the solution should remain in the container • Re-chill the remaining half if desired
<u>After Dose 1</u>	<ul style="list-style-type: none"> • Continue drinking Clear liquids • Drink at least 2 more cups (16oz) of clear liquids throughout the day
<u>Dose 2</u> Taken 6 hours prior to procedure time	<ul style="list-style-type: none"> • Drink the second half of the solution steadily over the next hour and a half <ul style="list-style-type: none"> • Drink 1 cup (8oz) every 10 minutes or so for a total of 8 cups (64oz) • The container should now be empty
<u>After Dose 2</u>	<ul style="list-style-type: none"> • Continue drinking Clear liquids up to 3 hours before procedure
<u>3 Hours Before</u>	<ul style="list-style-type: none"> • Nothing to eat or drink (other than essential medications) 3 hours before procedure

<u>Important</u>	
If you are taking any of these medications please contact the office ASAP	<ul style="list-style-type: none"> • Heparin, Lovenox, Warfarin (Coumadin), Rivaroxaban (Xarelto), Apixaban (Eliquis), Edoxaban (Lixiana, Savaysa), Betrixaban (Bevyxxa), Clopidigrel (Plavix), Prasugrel (Effient), Dabigatran (Pradaxa), Bivalirudin (Angiomax), Argatroban (Argatra, Novastan, Arganova, Exembol), and Desirudin (Iprivask, Revasc).
If you are diabetic	<ul style="list-style-type: none"> • Contact your Primary care doctor right away for instructions regarding your medications
If you are taking blood pressure medication	<ul style="list-style-type: none"> • DO NOT skip your blood pressure medication including the day before and day of your procedure
Female patients of childbearing age will be required to submit a urine sample for a pregnancy test upon arrival	

Please arrive at least 30 minutes prior to your appointment time
If you experience nausea, bloating or vomiting, try drinking the solution more slowly or cooling the solution on ice. Note: Individual responses to laxatives do vary; this prep may cause multiple bowel movements. This prep often works within 30 minutes, but it may take as long as 4-6 hours.
Someone must accompany you and take you home after the procedure. Do not drive for 8 hours after your procedure.
If you have any questions, please feel free to call your doctor’s office. Our phone numbers are listed above, and we will be happy to help you.

Please have your <u>COVID-19 PCR Test</u> completed 3-5 days prior to your procedure. (No Rapid test) If you have not had this test completed at a participating Northwell facility, please bring your results with you.
--

*** Please notify our office as soon as possible if you need to cancel your procedure.**
You will be charged a cancellation fee of \$200.00 if the procedure is not cancelled 48hrs before your scheduled procedure time.