

## Colonoscopy Instructions: Golytely

Time	Instructions		
<b><u>7 Days Before</u></b>	<ul style="list-style-type: none"> <li>• Stop all fiber supplements &amp; medications containing iron, including multivitamins and Omega 3</li> <li>• Stop taking Motrin, Advil &amp; Aleve; decrease Aspirin to 81mg daily</li> <li>• Be sure to drink 6-8 cups (8oz) of water each day leading up to your procedure</li> <li>• Be sure to have someone ready to pick you up after the procedure</li> <li>• Make sure you have your preparation medication (Golytely) by this time</li> </ul>		
<b><u>3 Days Before</u></b>	<table border="0" style="width: 100%;"> <tr> <td style="width: 50%; vertical-align: top;"> <u>Avoid High Fiber Foods including:</u> <ul style="list-style-type: none"> <li>• Raw Fruits &amp; Vegetables</li> <li>• Whole Wheat Bread/ Crackers</li> <li>• Seeds</li> <li>• Nuts</li> <li>• Popcorn, Bran</li> <li>• Quinoa</li> <li>• Corn</li> </ul> </td> <td style="width: 50%; vertical-align: top;"> <u>Begin a Low Fiber Diet such as:</u> <ul style="list-style-type: none"> <li>• White Bread &amp; Rice</li> <li>• Eggs</li> <li>• White Meat (Turkey, Chicken)</li> <li>• Fish</li> <li>• Cheese</li> <li>• Yogurt</li> <li>• Milk</li> <li>• Cooked Vegetables</li> </ul> </td> </tr> </table>	<u>Avoid High Fiber Foods including:</u> <ul style="list-style-type: none"> <li>• Raw Fruits &amp; Vegetables</li> <li>• Whole Wheat Bread/ Crackers</li> <li>• Seeds</li> <li>• Nuts</li> <li>• Popcorn, Bran</li> <li>• Quinoa</li> <li>• Corn</li> </ul>	<u>Begin a Low Fiber Diet such as:</u> <ul style="list-style-type: none"> <li>• White Bread &amp; Rice</li> <li>• Eggs</li> <li>• White Meat (Turkey, Chicken)</li> <li>• Fish</li> <li>• Cheese</li> <li>• Yogurt</li> <li>• Milk</li> <li>• Cooked Vegetables</li> </ul>
<u>Avoid High Fiber Foods including:</u> <ul style="list-style-type: none"> <li>• Raw Fruits &amp; Vegetables</li> <li>• Whole Wheat Bread/ Crackers</li> <li>• Seeds</li> <li>• Nuts</li> <li>• Popcorn, Bran</li> <li>• Quinoa</li> <li>• Corn</li> </ul>	<u>Begin a Low Fiber Diet such as:</u> <ul style="list-style-type: none"> <li>• White Bread &amp; Rice</li> <li>• Eggs</li> <li>• White Meat (Turkey, Chicken)</li> <li>• Fish</li> <li>• Cheese</li> <li>• Yogurt</li> <li>• Milk</li> <li>• Cooked Vegetables</li> </ul>		
<b><u>3-5 Days Before</u></b>	<ul style="list-style-type: none"> <li>• If <u>not</u> fully vaccinated: <u>get your mandatory COVID-19 PCR or Antigen test</u></li> <li>• If fully vaccinated (or have had COVID in the last three months): no COVID test required, but you are required to bring proof/documentation to your procedure</li> </ul>		
<b><u>24 Hours Before</u></b>	<ul style="list-style-type: none"> <li>• <b>NO</b> solid foods, <b>only</b> clear liquids including: <ul style="list-style-type: none"> <li>• Water</li> <li>• Apple, White Grape, &amp; White Cranberry juices without pulp</li> <li>• Clear soup Broth</li> <li>• Tea or Coffee (No milk, cream)</li> <li>• Gatorade / Powerade (<b>NO</b> red, orange, or purple colors)</li> <li>• Jell-O (<b>NO</b> red, orange, or purple colors)</li> <li>• Popsicles or Sorbet (<b>NO</b> red, orange, or purple colors)</li> <li>• No Alcohol</li> </ul> </li> </ul>		

<b><u>Important</u></b>	
If you are taking any of these medications please contact the office ASAP	<ul style="list-style-type: none"> <li>• Heparin, Lovenox, Warfarin (Coumadin), Rivaroxaban (Xarelto), Apixaban (Eliquis), Edoxaban (Lixiana, Savaysa), Betrixaban (Bevyxxa), Clopidigrel (Plavix), Prasugrel (Effient), Dabigatran (Pradaxa ), Bivalirudin (Angiomax), Argatroban (Argatra, Novastan, Arganova, Exembol), Brilinta (Ticagrelor) and Desirudin (Iprivask, Revasc).</li> </ul>
If you are diabetic	<ul style="list-style-type: none"> <li>• Contact your Primary care doctor right away for instructions regarding your medications</li> </ul>
If you are taking blood pressure medication	<ul style="list-style-type: none"> <li>• <b>DO NOT</b> skip your blood pressure medication including the day before and day of your procedure</li> </ul>
<b>Please arrive at least 30 minutes prior to your appointment time</b>	
Female patients of childbearing age will be required to submit a urine sample for a pregnancy test upon arrival	

**\* Please notify our office as soon as possible if you need to cancel your procedure. You will be charged a cancellation fee of \$200.00 if the procedure is not cancelled 48hrs before your scheduled procedure time.**



311 E 79 <sup>th</sup> St.	16 E 52 <sup>nd</sup> St.	68 E 86th St.	1150 Fifth Ave.	300 Cadman Plaza
212-996-6633	212-826-3903	212-535-1845	212-369-2490	718-280-0020

**Preparation Instructions: Golytely**

<b>Time</b>	<b>Instructions</b>
<b><u>24 Hours Before</u></b>	<ul style="list-style-type: none"> <li>• Prepare Golytely               <ul style="list-style-type: none"> <li>• Fill container to “fill” line with water and mix until dissolved</li> <li>• Chill if desired</li> </ul> </li> <li>• <b><u>NO solid foods, only clear liquids including:</u></b> <ul style="list-style-type: none"> <li>• Water</li> <li>• Apple, White Grape, &amp; White Cranberry juices without pulp</li> <li>• Clear soup Broth</li> <li>• Tea or Coffee (No milk, cream)</li> <li>• Gatorade / Powerade (<b>NO</b> red, orange, or purple colors)</li> <li>• Jell-O (<b>NO</b> red, orange, or purple colors)</li> <li>• Popsicles or Sorbet (<b>NO</b> red, orange, or purple colors)</li> <li>• No Alcohol</li> </ul> </li> </ul>
<b><u>Dose 1</u></b> Taken at 5pm night before procedure	<ul style="list-style-type: none"> <li>• Drink the first half of the Golytely solution steadily over the next hour and a half               <ul style="list-style-type: none"> <li>• Drink 1 cup (8oz) every 10 minutes or so for a total of 8 cups (64oz)</li> </ul> </li> <li>• Half of the solution should remain in the container</li> <li>• Re-chill the remaining half if desired</li> </ul>
<b><u>After Dose 1</u></b>	<ul style="list-style-type: none"> <li>• Continue drinking clear liquids</li> <li>• Drink at least 2 more cups (16oz) of clear liquids throughout the day</li> </ul>
<b><u>Dose 2</u></b> Taken 6 hours prior to procedure time	<ul style="list-style-type: none"> <li>• Drink the second half of the Golytely solution steadily over the next hour and a half               <ul style="list-style-type: none"> <li>• Drink 1 cup (8oz) every 10 minutes or so for a total of 8 cups (64oz)</li> </ul> </li> <li>• The container should now be empty</li> </ul>
<b><u>After Dose 2</u></b>	<ul style="list-style-type: none"> <li>• Continue drinking clear liquids up to 3 hours before procedure</li> </ul>
<b><u>3 Hours Before</u></b>	<ul style="list-style-type: none"> <li>• Nothing to eat or drink (other than essential medications) 3 hours before procedure</li> </ul>

**Important**

If you experience nausea, bloating or vomiting, try drinking the solution more slowly or cooling the liquid on ice.

Note: Individual responses to laxatives do vary; this prep may cause multiple bowel movements. This prep often works within 30 minutes, but it may take as long as 4-6 hours.

**Someone must accompany you and take you home after the procedure. Do not drive for 8 hours after your procedure.**

If you have any questions, please feel free to call your doctor’s office. Our phone numbers are listed above and we will be happy to help you.

If you are not fully vaccinated, you are required to take a COVID-19 test 3-5 days prior to your procedure. If you are fully vaccinated, or have had COVID in the last 3 months, you do not need a COVID test, but you must bring proof to your procedure. If you did not complete your test at a participating Northwell facility, please bring the results with you.